

Mindfulness and Self-Care for Behavioral Health Professionals

A Practical, Evidence-Based Guide

Prepared by: Ricc Terranova, LMHC — Broadlawns Medical Center, Des Moines, Iowa

Companion to: *Work, Life, and a Little Mindfulness*, IBC C.A.R.E. Conference¹

Written for: Substance use disorder counselors, mental health counselors, social workers, peer recovery specialists, and clinical supervisors

Reproduction: Permitted and encouraged for team meetings, supervision, and staff training

How to use this guide. Section 1 documents the occupational risk this workforce carries. Sections 2 and 3 explain why the risk is physiological rather than a matter of willpower. Section 4 summarizes the research evidence, including its weaknesses. Section 5 contains seven brief practices with full instructions. Sections 6 through 8 cover contraindications, organizational responsibility, and implementation. Readers who want only the practices can go straight to Section 5.

1. The Occupational Risk This Workforce Carries

Burnout in behavioral health is not an individual failing distributed at random. It is a predictable occupational outcome with measurable prevalence, and the data specific to substance use disorder treatment is more severe than the data for helping professions generally.

Exhaustion is close to the base rate

A survey of 699 addiction counselors across six states found emotional exhaustion averaging slightly above published norms for mental health providers. The more useful finding concerns causation: mindfulness and values-based psychological processes predicted burnout over and above workplace factors including workload, salary, supervisor support, and job control. The final models accounted for 27 percent of the variance in exhaustion.³

Two independent contributors therefore exist. Organizational conditions matter, and a clinician's relationship to their own internal experience matters separately. Interventions aimed at only one of the two will underperform.

Turnover is high and clinically consequential

- Average annual turnover across the behavioral health workforce is estimated at approximately 30 percent.⁵ In substance abuse treatment specifically, counselor turnover has been estimated near 32 percent annually and as high as 50 percent in some settings.⁶
- In one multi-site trial of SUD treatment, counselor turnover ranged from 33 to 74 percent across roughly two years — among counselors who had stated at enrollment that they had no intention of

leaving within that window.⁴

- Replacement of a trained clinician is commonly estimated at 1.5 to 2 times annual salary once recruitment, onboarding, lost productivity, and training are counted.^{5,6}

The clinical cost is separate from the financial one. Each departure interrupts continuity of care. A client in early recovery who has recounted their history to four counselors in eighteen months has received repeated intake rather than sustained treatment.

Vicarious trauma is an exposure, not a vulnerability

A 2025 multistate survey of opioid treatment program staff across 16 states found that 89.2 percent reported vicarious trauma exposure at work, 63.8 percent reported moderate-to-severe vicarious trauma, and 13 percent met preliminary criteria for PTSD.⁷ Framed as any other occupational exposure would be framed, prevalence at that level warrants systematic control measures rather than individual resilience training alone.

Existing self-care efforts are not sufficient on their own

In the American Psychological Association's 2023 Practitioner Pulse Survey, 36 percent of licensed psychologists reported feeling burned out despite engaging in self-care strategies. More than half reported increased symptom severity among their patients, and 56 percent reported no openings for new clients.⁸ Qualitative research with residential addiction treatment providers points in the same direction, identifying working conditions, understaffing, and low pay as the primary drivers of turnover.²¹

The interpretive question this guide addresses. If a third of practitioners are burned out while already practicing self-care, either self-care is ineffective or the activities commonly labeled self-care are not the activities that were studied. The evidence reviewed in Section 4 supports the second interpretation.

2. Why Sustained Overwork Degrades Clinical Work

The argument against work-life balance is usually made in economic terms. In clinical settings it fails on physiological grounds.

Cortisol and prefrontal function

Chronic overwork sustains cortisol elevation, and sustained cortisol impairs prefrontal functioning — the neural basis of decision-making, cognitive flexibility, and emotional regulation. In behavioral health these are not productivity variables. They are clinical judgment, impulse regulation within a session, and the capacity for empathic attunement.

Diminishing returns past roughly fifty hours

Productivity declines sharply beyond approximately 50 hours per week. A fatigued worker requires substantially longer to complete simple tasks and commits considerably more errors. In this field, error categories include a missed suicide screen, a rushed safety plan, and documentation that is never completed.

Unstructured rest performs cognitive work

During rest without external input, the brain engages the default mode network, which supports memory consolidation, processing of emotional material, and non-deliberate associative thinking. Unscheduled minutes between clients are therefore not idle time. They are when case formulation frequently consolidates.

3. Recognizing Burnout Before It Becomes a Crisis

Burnout rarely presents dramatically. The typical presentation is quiet withdrawal, hypervigilance, cynicism, and gradual disengagement, and it is routinely missed by supervisors because the affected clinician continues to meet obligations.

In behavioral health the earliest reliable indicator is usually depersonalization toward clients — the point at which a person becomes "the 2 o'clock," "a frequent flyer," or "a manipulator." This language is a measurable symptom with an established subscale score, not a personality characteristic. Supervisors who track how a team's language about clients changes over six months are observing burnout progression in real time, and at a stage where it remains a supervision matter rather than a human resources one.

Practical indicators worth monitoring: hardening language about clients; declining willingness to consult; increased documentation lag; reduced tolerance for ambiguity in case discussion; and the phrase "I'm fine" offered in place of an answer.

4. What the Research Shows

The evidence base for mindfulness-based interventions with clinicians is genuine but uneven. The findings below are presented with their limitations attached, because a professional audience is entitled to both.

Finding 1: Burnout reduction in physicians, durable at fifteen months

Krasner and colleagues delivered an eight-week intensive plus ten-month maintenance program in mindfulness meditation, self-awareness exercises, and narrative medicine to 70 primary care physicians. At 15 months, emotional exhaustion fell from 26.8 to 20.0 (effect size 0.62), depersonalization fell 2.5 points (0.45), personal accomplishment rose (0.44), mindfulness rose substantially (1.12), and total empathy rose 4.6 points (0.45).⁹

Limitation: this was a pre-post design without a randomized control group, and participants self-selected into a mindfulness course. The result is best characterized as promising and durable rather than established. Its clinical significance lies in the direction of the two key variables: exhaustion decreased while empathy increased.

Finding 2: Meta-analyses show moderate effects at low certainty

- A meta-analysis of mindfulness-based interventions with primary healthcare professionals (10 studies, 417 participants) found a pooled reduction of 5.89 points on the emotional exhaustion subscale of the Maslach Burnout Inventory.¹⁰

- A meta-analysis of 15 randomized trials involving 1,165 nurses found a moderate reduction in stress immediately post-intervention (SMD -0.81), still present at three-month follow-up (SMD -0.69), with improvements in emotional exhaustion and depersonalization. Anxiety and depression did not improve significantly.¹¹

Limitation: the nursing review graded its own certainty of evidence as very low, citing small samples, absence of blinding, self-report outcomes, and wait-list controls.¹¹ The defensible summary is that mindfulness produces a reliably moderate effect on stress and exhaustion — consequential in a field losing roughly a third of its workforce annually, and not a remedy for an unmanageable caseload.

Finding 3: Substantial gains in self-compassion

- A meta-analysis of 28 studies ($N = 1,020$) of mindfulness-based interventions with health care professionals and trainees found a medium-to-large increase in self-compassion (Hedges $g = 0.61$) that grew rather than decayed at follow-up ($g = 0.76$ at an average of 15 weeks). Control groups showed essentially no change ($g = 0.04$).¹²
- A larger meta-analysis of 56 randomized trials of self-compassion interventions found medium effects on depressive symptoms (SMD 0.44) and stress (SMD 0.43), and small-to-medium effects on anxiety (SMD 0.36), with an average intervention length of only 5.1 weeks.¹³

Self-compassion is the construct most directly relevant to career attrition in this field, because it targets the two cognitions that most reliably precede it: "I should have caught that" and "a competent counselor would not feel this way."

Limitation: effects were clear against wait-list controls but not consistently superior to active comparison conditions.¹³ Structured practice outperforms no practice; mindfulness cannot yet be said to outperform every alternative.

Finding 4: A single training day moved burnout in addiction counselors

Hayes and colleagues randomized 90 licensed and certified Nevada alcohol and drug counselors to one of three one-day, six-hour workshops: Acceptance and Commitment Training, multicultural training, or an educational control on drug biology. Overall burnout in the ACT group — emotional exhaustion plus depersonalization — improved significantly from pre- to post-workshop (17.87 to 15.20 , $p = .005$) and outperformed the control condition. At three-month follow-up, ACT also significantly outperformed control on stigmatizing attitudes toward people with substance use disorders, and reduced the believability of stigmatizing thoughts at both time points.¹⁵

This is the study most directly transferable to this workforce, and it establishes that a brief, single-session dose can produce measurable change.

Finding 5: Clinician practice appears in client outcomes

Grepmaier and colleagues conducted a randomized, double-blind trial in which 18 psychotherapists in training were randomized to practice Zen meditation or not, and the outcomes of their 124 inpatients were compared over nine weeks. Patients of the meditating therapists rated their therapy significantly higher on clarification and problem-solving perspectives, rated the overall therapeutic result higher, and showed greater symptom reduction on the SCL-90-R Global Severity Index and on somatization, anxiety, anger/hostility, obsessiveness, and social insecurity.¹⁶

The patients were unaware of the assignment. This is the strongest available basis for treating clinician self-regulation as a component of clinical competence rather than a personal matter. Related

work on mindful self-care in healthcare practitioners finds that relational and structural self-care dimensions are the strongest correlates of clinical empathy.²⁴

Finding 6: The same skill set is an established client intervention

Bowen and colleagues randomized 286 adults leaving intensive SUD treatment to Mindfulness-Based Relapse Prevention, standard cognitive-behavioral relapse prevention, or treatment as usual, with 12-month follow-up. At three months there were no significant group differences. At 12 months, MBRP participants reported fewer drug-use days (3.06 versus 4.63 for treatment as usual) and substantially fewer heavy-drinking days (1.44 versus 4.65); 7.8 percent of MBRP participants reported any heavy drinking at 12 months, compared with 20.0 percent of the treatment-as-usual group.¹⁷

MBRP already appears on many treatment menus. Staff are frequently the only group to whom the same skill set is not systematically offered.

Finding 7: The effective dose can be brief and app-delivered

Bostock and colleagues randomized 238 employees to eight weeks of a smartphone mindfulness application at 10 to 20 minutes daily, or to a wait-list. The intervention group showed improved well-being, reduced job strain, increased perceived job control and workplace social support, and reduced anxiety and depressive symptoms.¹⁴

Summary of the evidence. Mindfulness-based practice for clinicians produces moderate, replicated effects on stress and emotional exhaustion; larger and durable effects on self-compassion; at least one randomized double-blind trial in which therapist practice registered in patient outcomes; and measurable change from a single six-hour workshop with addiction counselors. It is not a substitute for adequate staffing, and it is the best-supported intervention an individual clinician directly controls.

5. Seven Brief Practices

Each practice below takes under five minutes and requires no equipment, application, or quiet room. Each is presented twice: as a self-care practice, and as a clinical tool usable with clients. All seven are invitations rather than requirements. Practitioners who feel worse during a practice should stop, open their eyes, and reorient to the room; that response is informative and is addressed in Section 6.

Practice 1 — Watching Thoughts Pass (1 to 3 minutes)

- **Concept.** The metaphor attributed to Ram Dass: thoughts are fish swimming in water, and awareness is the ocean. The fish will swim; the practice is not catching them on a hook. The aim is not an empty mind.
- **Instructions.** Sit without changing anything. For 60 seconds, observe thoughts as they arrive and depart, labeling each silently as a passing event rather than a task. Unpaid bill: a thought. Unwritten note: a thought. Remain the observer.
- **Clinical use.** This is urge surfing at an earlier stage. Clients who cannot yet tolerate a craving can often tolerate a thought about a craving, which makes this a useful entry point.

Practice 2 — Dropping Anchor (2 to 3 minutes)

- **Concept.** An Acceptance and Commitment Therapy grounding sequence for use during emotional escalation, structured as ACE: Acknowledge, Connect, Engage.
 - **Acknowledge.** Name what is present — shoulder tension, racing thought, urge to leave — without altering it.
 - **Connect.** Press the feet into the floor. Press the fingertips together. Shrug the shoulders and release.
 - **Engage.** Open the eyes. Identify three things visible and two audible, and return attention to the room.
- **Clinical use.** Safe to use with a client mid-escalation and in the ninety seconds after they leave. Because it is externally anchored, it is generally the appropriate first practice for trauma-exposed clients.

Practice 3 — 4-7-8 Breathing (1 to 3 minutes)

- **Concept.** Extending the exhale relative to the inhale shifts autonomic balance from sympathetic activation toward parasympathetic recovery.
- **Instructions.** Empty the lungs. Inhale quietly through the nose for a count of four. Hold for seven. Exhale audibly through the mouth for eight. Three cycles.
- **Contraindication.** Breath-focused practice can be activating for clients with panic disorder or a trauma history involving restriction or strangulation. If breath counting increases anxiety, substitute a grounding practice. The breath should never be the only available entry point.

Practice 4 — Cognitive Defusion (30 seconds to 3 minutes)

- **Concept.** Creating distance between a thought and the behavior it prompts, so that the thought is observed rather than obeyed.
- **Instructions.** Identify one self-limiting professional thought from the past week — for example, "I am not good enough for this caseload" or "I should have seen that relapse coming." Repeat it silently prefixed with "I am having the thought that..." Then add a second layer: "I notice I am having the thought that..." Observe the space this creates.
- **Clinical use, with evidence.** In the Nevada counselor trial, ACT training reduced both burnout and the believability of stigmatizing thoughts about clients.¹⁵ Defusion is the mechanism by which "this client is manipulative" becomes "I am having the thought that this client is manipulative, and I am tired." The first is a diagnostic assertion; the second is an observation of one's own state.

Practice 5 — The Self-Compassion Break (2 to 4 minutes)

- **Concept.** The central exercise of Mindful Self-Compassion, developed by Neff and Germer. In their randomized trial, participants showed significant increases in self-compassion, mindfulness, life satisfaction, and happiness, and significant decreases in depression, anxiety, and stress, with gains retained at six-month follow-up.¹⁸ Meta-analytic evidence across 56 randomized trials supports self-compassion interventions for stress and depressive symptoms.^{13,22}
- **Instructions.** Three statements, said internally, optionally with a hand placed over the heart:
 - **Mindfulness.** "This is hard right now." Name the difficulty without dramatizing or minimizing it.

- **Common humanity.** "I am not the only counselor who feels this. Difficulty with this work is part of this work."
- **Self-kindness.** "May I offer myself what I would offer a client who said this to me." Then state the actual sentence that would be offered to that client.
- **Note on the mechanism.** The instructive step is repeating verbatim what was said to a client who reported failure this week. The discomfort that accompanies it marks the gap between compassion dispensed and compassion accepted.
- **Expected difficulty.** Some practitioners feel worse initially, particularly those for whom self-criticism has long functioned as a motivational strategy. This reaction is described in the literature as backdraft. It indicates a need to shorten and slow the practice rather than to abandon it.

Practice 6 — The Three-Step Breathing Space, or Doorway Reset (3 minutes)

- **Concept.** The Three-Minute Breathing Space from Mindfulness-Based Cognitive Therapy, designed as a portable practice for ordinary conditions rather than a formal sitting.²³ Its structure is an hourglass: wide attention, narrow, wide again.
 - **Step 1 — Attend.** What thought, feeling, and body sensation are present? Name each in one word. Do not correct them.
 - **Step 2 — Narrow.** Gather attention onto the breath at one location: nostrils, chest, or abdomen.
 - **Step 3 — Expand.** Widen to the whole body in the chair — posture, face, hands — then carry that awareness into the next task.
- **Why this one suits counselors.** It fits in the interval between clients, including when most of that interval belongs to documentation. It is the practical difference between carrying a 10:00 appointment into an 11:00 appointment and leaving it where it belongs.

Practice 7 — Loving-Kindness for a Difficult Client (2 to 3 minutes)

- **Concept.** Loving-kindness practice directed specifically at depersonalization, the burnout dimension with the most direct effect on clients. Extending goodwill toward a difficult person is the established advanced step of this practice, and compassion-based training reliably increases positive affect and reduces the self-protective distance associated with poorer outcomes.
 - Bring to mind the client whose cancellation would be a relief. Do not attempt to correct the reaction.
 - Silently: *May you be safe. May you be free from suffering. May you find your way, whether or not I am the one who sees it.*
 - Then direct the same phrases inward: *May I be safe. May I be free from suffering. May I be at ease. May I be tired and still be competent.*
- **Realistic expectation.** The practice will feel artificial at first. The aim is not to manufacture an emotion but to rehearse a professional stance in which the client remains a person rather than a slot on a schedule.
- **Supervisory use.** "Which client do you dread?" reaches countertransference material faster than most structured forms.

Dosing. Total practice time for all seven is approximately twenty minutes. Sustained benefit does not require an eight-week program. The app trial that produced measurable change used ten minutes daily;¹⁴ the workshop that moved burnout in addiction counselors lasted six hours.¹⁵ Selecting one practice and attaching it to an existing cue is sufficient to begin.

6. Contraindications, Adverse Effects, and Screening

Mindfulness is an intervention, and interventions have adverse effects. Presenting it otherwise undermines its credibility with clinicians and risks harm to clients.

- In an international survey of 1,370 regular meditators, 22 percent reported unpleasant meditation-related experiences and 13 percent reported experiences classified as adverse, predominantly affective, somatic, and cognitive.¹⁹
- A systematic review of 83 studies found an overall adverse-event prevalence of 8.3 percent, ranging from 3.7 percent in experimental studies to 33.2 percent in observational studies.²⁰

Practical screening guidance

- Exercise caution with active psychosis, unprocessed trauma, dissociative presentations, and severe depression with prominent rumination.
- Prefer eyes-open, externally anchored practices — grounding, walking, sensory orientation — over extended silent internal attention when working with trauma-exposed clients.
- Titrate duration upward from very short exposures, and debrief afterward as with any other intervention.
- Treat a worsening response as clinical information warranting discussion in supervision or personal therapy, not as evidence of insufficient effort.

7. Organizational Responsibility

Individual practice does not discharge the employer's obligation, and this guide should not be used to suggest otherwise. The evidence supports two conclusions simultaneously: workplace conditions and individual psychological flexibility independently predicted burnout among 699 addiction counselors,³ and qualitative research with residential addiction treatment providers identifies working conditions, staffing, and compensation as the drivers of turnover.²¹ SAMHSA's own guidance on behavioral health burnout is organized around organizational strategies rather than individual resilience.⁵

Recommendations for supervisors and administrators

- **Model the practice or omit the topic.** Messages sent at 11:00 PM communicate an expectation of availability regardless of stated policy. Delayed delivery is a low-cost correction. A leader's calendar functions as the operative wellness policy.
- **Change the check-in question.** Replace "where are you on documentation?" with "what is your current bandwidth, and what can come off your plate this week?"

- **Track depersonalization as a supervision metric.** Language about clients is a free, continuously available, early burnout indicator.
- **Schedule recovery time.** Ten unscheduled minutes between clients is a clinical intervention with a measurable return, not lost productivity, particularly when weighed against replacement costs of 1.5 to 2 times salary.^{5,6}
- **Address the exposure, not only the coping.** Where 89 percent of staff report vicarious trauma exposure, that exposure belongs in the organizational risk register alongside other occupational hazards.⁷

A distinction worth stating plainly. Distributing a meditation application in response to an unmanageable caseload transfers the cost of understaffing onto the clinician. Individual practice and organizational reform are complementary obligations, not substitutes.

8. Implementation: The Next Two Weeks

- **Select one practice.** Not seven. The most transferable choices for clinical schedules are Practice 6 (Doorway Reset) and Practice 2 (Dropping Anchor).
- **Attach it to an existing cue.** A hand on the door handle, sitting down in the car, or the moment a chart closes. Practices anchored to existing cues persist; practices dependent on recall do not.
- **Set a floor, not a target.** One three-minute practice, five days a week, is a defensible minimum. If that proves impossible, the constraint is caseload rather than technique, and the appropriate response is a workload conversation.
- **Secure a place to put the material.** Supervision is not therapy and consultation is not support. Clinicians who carry trauma content professionally require a dedicated outlet for it.
- **Review at two weeks.** Retain what was actually performed and discard what was not, rather than resolving to try harder.

The core proposition. In behavioral health the practitioner is the instrument of treatment. A clinician's regulation is therefore a clinical variable, not a personal amenity — and in at least one randomized double-blind trial, therapist practice was reflected directly in patient outcomes.¹⁶

Quick Reference: The Seven Practices

- **Watching Thoughts Pass (1 min).** Observe thoughts as passing events. No emptying of the mind required.
- **Dropping Anchor / ACE (2 min).** Acknowledge what is present. Connect with the body. Engage with the room.
- **4-7-8 Breathing (1 min).** Inhale 4, hold 7, exhale 8. Three cycles. Omit if breath focus increases anxiety.

- **Cognitive Defusion (30 sec).** "I am having the thought that..." then "I notice I am having the thought that..."
- **Self-Compassion Break (2 min).** This is hard right now. I am not the only one. May I offer myself what I would offer a client.
- **Doorway Reset (3 min).** Attend to what is present, narrow to the breath, expand to the whole body. Use between clients.
- **Loving-Kindness for a Difficult Client (2 min).** May you be safe. May you be free from suffering. May you find your way. Then direct it inward.

Ricc Terranova, LMHC

Broadlawns Medical Center | Johnston & Des Moines, Iowa

Taking referrals and accepting new clients.

terranogetherapy515@hushmail.com

terranova-counseling.pplx.app



My practice



Email me

References

Full citations with links. Numbering matches the conference session materials.

1. Iowa Board of Certification, CARE 2026 Conference <https://iowabc.org/care-2026/>
2. Iowa Board of Certification, C.A.R.E. Conference call for presentations (C.A.R.E. pillars) <https://iowabc.org/2025-care-conference/>
3. Vilardaga R, Luoma JB, Hayes SC, et al. Burnout among the addiction counseling workforce: the differential roles of mindfulness and values-based processes and work-site factors. *J Subst Abuse Treat*. 2011;40(4):323-335. <https://pmc.ncbi.nlm.nih.gov/articles/PMC3081943/>
4. Hatch-Maillette MA, Harwick R, Baer JS, et al. Counselor turnover in substance use disorder treatment research: observations from one multi-site trial. *Subst Abus*. 2019;40(2):214-220. <https://pmc.ncbi.nlm.nih.gov/articles/PMC6759413/>
5. SAMHSA. Addressing Burnout in the Behavioral Health Workforce Through Organizational Strategies (PEP22-06-02-005). <https://library.samhsa.gov/sites/default/files/pep22-06-02-005.pdf>
6. SAMHSA. Quality Assurance Practices in Substance Abuse Treatment Facilities (Short Report 3059). https://www.samhsa.gov/data/sites/default/files/report_3059/ShortReport-3059.html
7. Coffee Z, Linde-Krieger L, Carter GA, et al. Trauma-related stress and resilience in a multistate sample of methadone treatment staff. *Subst Use Res Treat*. 2025;19. <https://pmc.ncbi.nlm.nih.gov/articles/PMC12536211/>
8. American Psychological Association. 2023 Practitioner Pulse Survey: Psychologists reaching their limits. <https://www.apa.org/pubs/reports/practitioner/2023-psychologist-reach-limits>
9. Krasner MS, Epstein RM, Beckman H, et al. Association of an educational program in mindful communication with burnout, empathy, and attitudes among primary care physicians. *JAMA*. 2009;302(12):1284-1293. <https://jamanetwork.com/journals/jama/fullarticle/184621>
10. Salvado M, Marques DL, Pires IM, Silva NM. Mindfulness-based interventions to reduce burnout in primary healthcare professionals: a systematic review and meta-analysis. *Healthcare*. 2021;9(10):1342. <https://pmc.ncbi.nlm.nih.gov/articles/PMC8544467/>
11. Wang Q, Wang F, Zhang S, et al. Effects of mindfulness-based interventions on stress and burnout in nurses: a systematic review and meta-analysis. *Front Psychiatry*. 2023;14:1218340. <https://www.frontiersin.org/journals/psychiatry/articles/10.3389/fpsy.2023.1218340/full>
12. Wasson RS, Barratt C, O'Brien WH. Effects of mindfulness-based interventions on self-compassion in health care professionals: a meta-analysis. *Mindfulness*. 2020;11(8):1914-1934. <https://pmc.ncbi.nlm.nih.gov/articles/PMC7223423/>
13. Han A, Kim TH. Effects of self-compassion interventions on reducing depressive symptoms, anxiety, and stress: a meta-analysis. *Mindfulness*. 2023;14. <https://pmc.ncbi.nlm.nih.gov/articles/PMC10239723/>
14. Bostock S, Crosswell AD, Prather AA, Steptoe A. Mindfulness on-the-go: effects of a mindfulness meditation app on work stress and well-being. *J Occup Health Psychol*. 2019;24(1):127-138. <https://www.apa.org/pubs/journals/features/ocp-ocp0000118.pdf>
15. Hayes SC, Bissett R, Roget N, et al. The impact of acceptance and commitment training and multicultural training on the stigmatizing attitudes and professional burnout of substance abuse counselors. *Behav Ther*. 2004;35(4):821-835. <https://core.ac.uk/download/pdf/214049460.pdf>
16. Grepmaier L, Mitterlehner F, Loew T, et al. Promoting mindfulness in psychotherapists in training influences the treatment results of their patients: a randomized, double-blind, controlled study. *Psychother Psychosom*. 2007;76(6):332-338. <https://pubmed.ncbi.nlm.nih.gov/17917468/>

17. Bowen S, Witkiewitz K, Clifasefi SL, et al. Relative efficacy of mindfulness-based relapse prevention, standard relapse prevention, and treatment as usual for substance use disorders: a randomized clinical trial. *JAMA Psychiatry*. 2014;71(5):547-556. <https://pmc.ncbi.nlm.nih.gov/articles/PMC4489711/>
18. Neff KD, Germer CK. A pilot study and randomized controlled trial of the Mindful Self-Compassion program. *J Clin Psychol*. 2013;69(1):28-44. <https://self-compassion.org/wp-content/uploads/2014/10/Neff-Germer-MS-C-RCT-2012.pdf>
19. Pauly L, Bergmann N, Hahne I, et al. Prevalence, predictors and types of unpleasant and adverse effects of meditation in regular meditators: international cross-sectional study. *BJPsych Open*. 2021;8(1):e11. <https://pmc.ncbi.nlm.nih.gov/articles/PMC8693904/>
20. Farias M, Maraldi E, Wallenkampf KC, Lucchetti G. Adverse events in meditation practices and meditation-based therapies: a systematic review. *Acta Psychiatr Scand*. 2020;142(5):374-393. <https://onlinelibrary.wiley.com/doi/10.1111/acps.13225>
21. Stelson EA, Sabbath-Clayton LL, Sorensen G, Sabbath EL. Working Conditions, Worker Health and Wellbeing, and Turnover of Residential Addiction Treatment Providers in Massachusetts. Harvard T.H. Chan School of Public Health. https://hsph.harvard.edu/wp-content/uploads/2025/09/SUP_wellbeing_study_report.pdf
22. Greater Good in Action, UC Berkeley. Self-Compassion Break (practice instructions). https://ggia.berkeley.edu/practice/self_compassion_break
23. Mindful.org. The Three-Minute Breathing Space Practice (Segal, Williams and Teasdale, MBCT). <https://www.mindful.org/the-three-minute-breathing-space-practice/>
24. Mindful self-care and clinical empathy in healthcare practitioners (2025). <https://pmc.ncbi.nlm.nih.gov/articles/PMC13097769/>